http://dx.doi.org/10.1590/1516-4446-2019-0583
Suicide and suicide attempts in adults: exploring suicide risk 24 months after a psychiatric emergency room visit - Taron M et al.

Online-only supplementary material

Table S1 Sample characterization (n=147)

Variables	Total, n (%)
Sociodemographic	
Gender	
Male	56 (38.1)
Female	91 (61.9)
Age	
Mean (±SD); min-max (years)	41.93 (±11.9); 19-65
18-24	13 (8.8)
25-34	29 (19.7)
35-44	43 (29.3)
45-54	35 (23.8)
55-65	27 (18.4)
Marital status	
Married	59 (40.1)
Others	88 (59.9)
Employment	
Active	58 (39.5)
Inactive	89 (60.5)
Clinical	
Personality disorders	
No	60 (40.8)
Yes	87 (59.2)
Borderline	40 (27.2)
Others	47 (32.0)
Psychiatric diagnoses	
No	8 (5.4)
Yes	139 (94.6)
Mood disorders	85 (57.8)
Alcoholic disorders	14 (9.5)
Psychotic disorders	14 (9.5)
Anxiety disorders	5 (3.4)
Others	21 (14.3)
Family history of psychiatric pathology	
No	102 (69.4)
Yes	45 (30.6)
Family history of suicide attempts	
No	133 (90.5)
Yes	14 (9.5)
Family history of suicide consummations	
No	118 (80.3)
Yes	29 (19.7)

Impulsivity	
No	28 (19.0)
Yes	119 (81.0)
Suicidal ideation	
No	7 (4.8)
Yes	140 (95.2)
NSSI	
No	105 (71.4)
Yes	42 (28.6)
Cuts	30 (20.4)
Others	12 (8.2)
Previous suicide attempts	
Nº of previous SA (n= 147)	
Mean (±SD); min-max	1.93 (±1.83); 0-10
0	30 (20.4)
1	47 (32.0)
≥ 2	70 (47.6)
Lethal method	
Drug overdose	84 (72.4)
Other	32 (27.6)
Health services utilization	
Nº of previous psychiatric emergencies	
Mean (±SD); min-max	2.59 (±4.51); 0-34
0	55 (37.4)
1	28 (19.0)
2	14 (19.5)
≥ 3	50 (34.0)
Nº of previous psychiatric admissions	
Mean (±SD); min-max	1.12 (±2.12); 0-10
0	23 (15.6)
1	89 (60.5)
2	16 (10.9)
≥ 3	19 (12.9)
Time since the last psychiatric hospital admission	
< 1 year	24 (16.3)
≥ 1 year	34 (23.1)

min-max = minimum-maximum; NSSI = non-suicidal self-injuries; SA = suicide attempt; SD = standard deviation.
* Due to missing values, we present only valid percentages.